Weill Cornell Imaging



WCINYP Access Center: 212.746.6000 Fax: 646.962.0122

wcinyp.com

Patient Information *Required Information			
*Patient Name		*DOBN	IYPH / MRN
*Signs and Symptoms/ICD-10 Codes			
Appointment Information		Patients: Please review exam pr	reparation instructions on reverse side
	Exam Time P		
Location: ☐ 1305 York Ave ☐ 1283 York Ave ☐ 520 E 70th St ☐ 425 E 61st St ☐ 416 E 55th St ☐ 2315 Broadway ☐ 53 Beekman St ☐ 28-25 Jackson Ave			
Physician Information			se fax completed form to 646.962.0122
	lo .lop (:		
STAT Read: Call () report Send CD of images: to office with patient report only Reports are mailed/faxed automatically			
MRI □ w/o contrast □ w/wo contrast	CT	PET/MRI	BREAST IMAGING
☐ 3D Recon. ☐ Arthrography (MSK exams)☐ Radiologist may change contrast	☐ w/ contrast ☐ w/wo contrast ☐ w/o contrast ☐ 3D Recon. ☐ Radiologist may change contrast	Specify both PET and MRI below: PET	☐ Radiologist may complete workup as clinically indicated ☐ Screening Mammography
order if clinically indicated	order if clinically indicated Head	☐ Brain [78608] ☐ Skull Base to Mid-Thigh [78812]	Includes 3D Tomosynthesis
☐ Brain ☐ Orbits	☐ Sinuses	☐ Whole Body (top of head to toes) [78813]	☐ Diagnostic Mammography ☐ Bilateral ☐ Right ☐ Left
□ IAC □ Pituitary	☐ Orbits ☐ Temporal Bone	MRI	☐ Breast Ultrasound ☐ Bilateral ☐ Right ☐ Left
☐ TMJ☐ Soft Tissue Neck	☐ Facial Bones ☐ Nasopharynx	□ w/o contrast □ w/wo contrast □ Radiologist may change contrast	│
□ Nasopharynx □ Cervical Spine	☐ TMJ for Prostheses Fitting ☐ Neck	order if clinically indicated	□ Ductograms □ Right □ Left
☐ Thoracic Spine	☐ Chest	☐ Brain ☐ Orbit, Face and/or Neck	☐ MRI-guided Biopsy ☐ Right ☐ Left
☐ Lumbar Spine ☐ T2- Iron Overload	☐ Lung Screening ☐ Pulmonary Embolism Study	☐ Abdomen	☐ Ultrasound-guided FNA
☐ Abdomen ☐ Heart ☐ Cardiac	☐ Cardiac	☐ Pelvis ☐ Breast	│
☐ Adenosine Perfusion	☐ Congenital ☐ PVI ☐ Non-coronary ☐ Calcium Scoring Only ☐ TAVR	☐ Other	☐ Right ☐ Left ☐ Other
☐ Chest ☐ Breast	☐ Abdomen and Pelvis ☐ Abdomen Only	ULTRASOUND	BONE DENSITY (DEXA)
☐ Abdomen	☐ Pelvis Only ☐ Enterography	☐ add Doppler if indicated	☐ Hip and/or Lumbar Spine
□ Elastography □ Brachial Plexus	☐ Urogram	☐ Abdominal ☐ Complete	☐ FRAX ☐ Trabecular Bone Score
☐ MRCP ☐ with Secretin / CCK	☐ Colonography (Virtual Colonoscopy)☐ Cervical Spine	☐ Limited (e.g. RUQ, LUQ, Ascites,	☐ Vertebral Fracture Assessment ☐ Appendicular Skeleton
☐ Pelvis	☐ Thoracic Spine	Spleen, Appendix, Hernia) Pelvic Transabdominal	☐ Body Composition
☐ Enterography ☐ Defecography	☐ Lumbar Spine	☐ Add transvaginal if indicated ☐ Pelvic Transvaginal	GENERAL X-RAY
□ Prostate □ Shoulder □ Right □ Left	☐ Extremity ☐ Other	☐ Obstetrics (<14 weeks)	☐ Chest: # of views
□ Hip □ Right □ Left	CT ANGIOGRAPHY	☐ Infant Hips ☐ Infant Head	☐ Cervical Spine
☐ Knee ☐ Right ☐ Left ☐ Ankle ☐ Right ☐ Left	☐ Cardiac	☐ Bladder	☐ Thoracic Spine ☐ Lumbar Spine
☐ Foot ☐ Right ☐ Left	☐ w/ Calcium Score ☐ TAVR ☐ Add FFR _{CT} if indicated	☐ Renal ☐ Renal (for Renal Artery Stenosis)	☐ Sacrum/Coccyx
☐ Other	☐ Head ☐ Neck	☐ Renal Transplant ☐ Liver Transplant	☐ Scoliosis☐ Osseous Survey
MR ANGIOGRAPHY	☐ Chest ☐ Abdomen ☐ Pelvis ☐ Lower Extremity	☐ Scrotal (with Doppler)	☐ Pelvis☐ Abdomen
☐ Head ☐ Neck (Carotid)	☐ Other	☐ Thyroid☐ Aorta	☐ Series ☐ KUB
☐ Chest`☐ Abdomen	PET/CT	☐ Screening ☐ Diagnostic	□ Ribs □ Right □ Left
⊐ Pelvis	☐ Skull Base to Mid-Thigh	☐ Hysterosonography	│ ☐ Extremity: ☐ Right ☐ Left
☐ Extremities ☐ Lower ☐ Upper	with Vertex	Soft Tissue (e.g. neck, lymph nodes, palpable mass)	Other
☐ Perforated Flap (PFA) ☐ Abdomen / Pelvis	☐ Whole Body (top of head to toes) ☐ Melanoma or Multiple Myeloma	☐ Extremities Non-vascular (MSK) ☐ Upper ☐ Lower (e.g. Baker's cyst)	FLUOROSCOPY
☐ Pelvis / Lower Extremity	Other indication	☐ Right ☐ Left	☐ Esophagram
☐ Other	☐ Brain ☐ Dementia ☐ Seizure ☐ Tumor	☐ Other	☐ Modified Barium Swallow ☐ Upper GI
MR VENOGRAPHY	☐ Other indication	VASCULAR	□ w/ Small Bowel
☐ Head ☐ Abdomen	☐ Limited Area PET/CT	☐ Carotid☐ Venous Doppler (r/o DVT)	☐ Small Bowel ☐ Enema
☐ Pelvis☐ Extremities	(specify)	□ Lower Extremity □ Upper Extremity	☐ Cystogram ☐ Tube Study
☐ Lower ☐ Upper	☐ Sarcoid/Inflammation ☐ Rest/Stress	☐ Right ☐ Left ☐ Bilateral	☐ Voiding Cystourethrogram (VCUG)
¬ Other	☐ Other	☐ Other	∏ Other

Patient Checklist

If you are pregnant, may be pregnant, or on a fertility protocol, please notify the technologist before the exam.

- ☐ Bring this prescription form to your appointment
- Consult your insurance company for any inquiries regarding referrals and pre-authorizations
- ☐ Bring prior outside CD images to your exam for comparison
- ☐ Bring your insurance card to your appointment (copays are collected at the time of service)
- ☐ If you are interested in receiving your results electronically, ask our staff about Weill Cornell Connect

Weill Cornell Imaging at NewYork-Presbyterian Locations



- **1** 1305 York Ave, 3rd Floor, NY, NY 10021 (70th St)
- **1283 York Ave**, 7th Floor, NY, NY 10065 (69th St)
- 🛘 🗿 **520 East 70th St**, Starr Pavilion, Floor 0, NY, NY 10021
- **425 East 61st St**, 9th Floor, NY, NY 10065
- **1 5 416 East 55th St**, Ground Floor, NY, NY 10022
- 2315 Broadway, 4th Floor, NY, NY 10024 (84th St)
- 53 Beekman St, Ground Floor, NY, NY 10038
- **3 28-25 Jackson Ave**, 2nd Floor, Long Island City, NY 11101

Exam Preparations

BONE DENSITY

- You may take calcium supplements and medications as prescribed.
- If coordinating with nuclear scans or IV contrast exams, the bone density scan must be done first.

CT

- Intravenous (IV) Contrast and Oral Contrast exams:
- IV Contrast: You will receive an injection of lodine-based dye at the time of the exam.
- Oral Contrast: You will be given oral contrast to drink 90 minutes prior to the exam. Length of stay is a minimum of 2 hours.
- Abdomen with Oral Contrast: Length of stay is a minimum of 1.5 hours (2 hours for Abdomen and Pelvis with Oral Contrast).
- Cardiac CTA: No Phosphodiesterase Type 5 Inhibitor such as VIAGRA®, CIALIS®, LEVITRA®, etc.) for 2 days prior to the exam.
- Colonography (Virtual Colonoscopy): Carefully follow the instructions regarding fasting, clear fluids and the bowel cleansing prep prescribed by your physician. You must pick up prep at Weill Cornell Imaging, at least 2 days prior to the exam.
- Patients with Diabetes or Decreased Kidney Function: It is recommended that BUN and creatinine blood levels be obtained by your physician and results be forwarded to Weill Cornell Imaging prior to the exam (lab work must be done within 2 weeks of exam).
- Diabetic Patients: If on an oral hypoglycemic such as Glucophage[®], you should stop these agents for 48 hours after the exam if IV contrast was received.

FLUOROSCOPY

- Esophagram, Small Bowl, and Upper GI: No food or liquids 8 hours prior to the exam.
- Modified Barium Swallow, Enema: Fasting requirements vary.
 Contact us for preparation information.

MAMMOGRAPHY

- Do not use deodorant, talcum powder, lotion or oils on day of exam.
- Notify the staff ahead of time if you have breast implants.
- Send prior outside films for comparison prior to appointment. For questions, please call 646.962.9617.

MR

- METAL Implants: Inform your referring physician and the MRI staff if you have any metal in your body including a CARDIAC PACEMAKER or IMPLANTABLE CARDIOVERTER-DEFIBRILLATOR (ICD)
- Abdomen: No food or liquids 2 hours prior to the exam.
- Pelvis: No food or liquids 2 hours prior to the exam. For gynecologic related exams, no food or liquids 6 hours prior to the exam.
- Head/Orbits/Neck/Face: Avoid wearing dry shampoo and any eye make-up or mascara during the exam.
- Patients with Decreased Kidney Function: Inform the MRI staff at the time of visit.

PET/CT and PET/MRI

- No food 4 hours prior to the exam (no insulin during this time).
- Do not eat candy or chew gum 4 hours prior to the exam.
- No sugar-containing drinks 4 hours prior to exam. Do not drink sodas, juices, shakes of any kind; no iced tea, coffee or tea if they contain sugar. You may drink water.
- · No exercise for 24 hours prior to the exam.
- Length of stay is a minimum of 2.5 hours.
- Diabetic Patients: Inform scheduler when making appointment in order to coordinate medication and diet prior to the exam. Blood sugar levels should be under control prior to the exam.

ULTRASOUND

- Abdomen: No food or liquids 6 hours prior to the exam. You may have water, and coffee or tea without milk or sugar. Morning appointments are preferred. If medication must be taken on the day of the exam, you may do so with water only.
- Pelvic and Renal: You should arrive with a full bladder (32 oz. of water is suggested), except for pelvic transvaginal and renal transplant exams. You should not urinate prior to the exam.
- Renal with Abdomen: Follow Abdomen preparation above.